arXiv:1303.4630v2 [math.AG] 10 Apr 2013

ON THE FUNDAMENTAL DOMAIN OF AFFINE SPRINGER FIBERS

ZONGBIN CHEN

Abstract. For G a connected reductive group, $\gamma \in \mathfrak{g}(F)$ semisimple regular unramified, we introduce a fundamental domain F_γ for the affine Springer fibers $\mathscr{X}_\gamma$. There is a beautiful way to reduce the purity conjecture of $\mathscr{X}_\gamma$ to that of F_γ, we call it the Arthur-Kottwitz reduction. It turns out that these fundamental domains behave well in family. We formulate a rationality conjecture about a generating series of their Poincaré polynomials. We then study them in detail for the group GL_3. In particular, we pave them in affine spaces and we prove the rationality conjecture.

1. Introduction

Let $k = \mathbf{F}_q$, we fix an algebraic closure $\bar{k}$ of k. Let $F = k((\epsilon))$ be the field of Laurent series with coefficients in k, $\mathcal{O} = k[[\epsilon]]$ the ring of integers of F, $\mathfrak{p} = \epsilon k[[\epsilon]]$ the maximal ideal of $\mathcal{O}$. Let $\mathrm{val} : F^\times \to \mathbf{Z}$ be the discrete valuation normalized by $\mathrm{val}(\epsilon) = 1$.

Let G be a connected reductive algebraic group split over k, we make the assumption that $\mathrm{char}(k) > \mathrm{rk}(G)$, where $\mathrm{rk}(G)$ is the semisimple rank of G. Let T be a split maximal torus of G, let B be a Borel subgroup of G containing T. Their Lie algebras will be denoted by the corresponding Gothic letters. Let $K = G(\mathcal{O})$ be the standard maximal compact subgroup of $G(F)$. The affine grassmannian $\mathscr{X} = G(F)/K$ is the ind-k-scheme such that

$$\mathscr{X}(\mathbf{F}_{q^n}) = G(\mathbf{F}_{q^n}((\epsilon)))/G(\mathbf{F}_{q^n}[[\epsilon]]), \quad \forall\, n \in \mathbf{N}.$$

For a regular element $\gamma \in \mathfrak{t}(\mathcal{O})$, the affine Springer fiber $\mathscr{X}_\gamma$ at γ

$$\mathscr{X}_\gamma = \{g \in G(F)/K \mid \mathrm{Ad}(g^{-1})\gamma \in \mathfrak{g}(\mathcal{O})\}$$

has been introduced by Kazhdan and Lusztig [KL]. It has been used by Goresky, Kottwitz and Macpherson [GKM1] to prove the fundamental lemma of Langlands-Shelstad in the unramified case, under the following hypothesis:

Conjecture 1.1 (Goresky-Kottwitz-Macpherson)**.** The cohomology of $\mathscr{X}_\gamma$ is pure in the sense of Deligne, i.e. the eigenvalues of the action of frobenius Fr_q on $H^i(\mathscr{X}_{\gamma,\bar{k}}, \overline{\mathbf{Q}}_l)$ have an absolute value of $q^{i/2}$ for any embedding $\overline{\mathbf{Q}}_l \to \mathbf{C}$.

This conjecture has been proved in some particular cases, see [GKM2], [Lu], [C], where the authors have found affine pavings of $\mathscr{X}_\gamma$.

The affine Springer fibers have a large symmetry group. The free abelian group Λ generated by $\chi(\epsilon)$, $\chi \in X_*(T)$ acts simply and transitively on the irreducible components of $\mathscr{X}_\gamma$. It is desirable to use this symmetry to reduce the study of $\mathscr{X}_\gamma$ to that of its irreducible components. But the condition of irreducibility is difficult to explore. Instead, we construct a fundamental domain F_γ of $\mathscr{X}_\gamma$ with respect to the action of Λ, which should be exactly one of the irreducible components of $\mathscr{X}_\gamma$. Our first main result is the following:

Theorem 1.1. *For any $\gamma \in \mathfrak{t}(\mathcal{O})$, suppose that F_γ^M is cohomologically pure for any Levi subgroup M of G containing T. Then $\mathscr{X}_\gamma$ is cohomologically pure if and only if F_γ is.*

The conjecture of Goresky, Kottwitz and Macpherson can be restated as:

Conjecture 1.2. The cohomology of F_γ is pure in the sense of Deligne.

It is believed that F_γ is an irreducible component of $\mathscr{X}_\gamma$, and that it is also the normalization of $\Lambda\backslash\mathscr{X}_\gamma$. For example, this follows from the purity conjecture. These observations lead us to restate the conjectures of Goresky, Kottwitz, Macpherson and of Laumon [L] in the following way, which opens the door to a possible proof of the purity conjecture by deformation.

Conjecture 1.3. Let C be a projective geometrically integral algebraic curve over k. Suppose that $\underline{\text{all}}$ the singularities of C are planar. Then the normalization of the compactified Jacobian $\overline{\mathrm{Jac}}_C$ of C is cohomologically pure.

Now we restrict to the group $G = \mathrm{GL}_d$. Let T be the maximal torus of diagonal matrices, let B be the Borel subgroup of G of the upper triangular matrices. Let $\Phi = \{\alpha_{i,j}\}$ be the root system of G with respect to T, let $\alpha_i = \alpha_{i,i+1}$, $i = 1, \cdots, d-1$, be the simple roots with respect to B. Let $\gamma \in \mathfrak{t}(\mathcal{O})$ be regular, it is said to be in minimal form if

$$\mathrm{val}(\alpha_{i,j}(\gamma)) = \min_{i\leq l\leq j-1}\{\mathrm{val}(\alpha_l(\gamma))\}, \quad \forall\, i<j.$$

In this case, we say that the root valuation of γ is $(\mathrm{val}(\alpha_1(\gamma)), \cdots, \mathrm{val}(\alpha_{d-1}(\gamma)))$. We can always conjugate γ such that it is in minimal form. For $\mathbf{n} = (n_1, \cdots, n_{d-1}) \in \mathbf{N}^{d-1}$, we can find $\gamma \in \mathfrak{t}(\mathcal{O})$ in minimal form with root valuation $\mathbf{n}$. It is believed that the topology of F_γ only depends on its root valuation. Let $P_{\mathbf{n}}(t)$ be its Poincaré polynomial.

Conjecture 1.4. The power series

$$\sum_{n_1=1}^{+\infty}\cdots\sum_{n_{d-1}=1}^{+\infty} P_{(n_1,\cdots,n_{d-1})}(t)\, T_1^{n_1}\cdots T_{d-1}^{n_{d-1}} \in \mathbf{Z}[[t, T_1, \cdots, T_{d-1}]]$$

is a rational function, i.e. it is an element of $\mathbf{Z}(t, T_1, \cdots, T_{d-1})$.

The second main result of our article is the following:

Theorem 1.2. *Let $G = \mathrm{GL}_3$, let $\mathbf{n} = (n_1, n_2) \in \mathbf{N}^2$, $n_1 \leq n_2$, let $\gamma \in \mathfrak{t}(\mathcal{O})$ be in minimal form with root valuation $\mathbf{n}$. The fundamental domain F_γ can be paved in affine spaces, and the paving only depends on $\mathbf{n}$. Its Poincaré polynomial is*

$$P_{\mathbf{n}}(t) = \sum_{i=1}^{n_1} i(t^{4i-2}+t^{4i-4}) + \sum_{i=2n_1}^{n_1+n_2-1}(2n_1+1)t^{2i} + \sum_{i=n_1+n_2}^{2n_1+n_2-1} 4(2n_1+n_2-i)t^{2i} + t^{4n_1+2n_2}.$$

The rationality conjecture for GL_3 is an easy consequence of the theorem.

Notations. Let $\Phi = \Phi(G,T)$ be the root system of G with respect to T, let W be the Weyl group of G with respect to T. For any subgroup H of G which is stable under the conjugation of T, we note $\Phi(H,T)$ for the roots appearing in $\mathrm{Lie}(H)$. Let Δ_B be the set of simple roots with respect to B, let $(\varpi_\alpha)_{\alpha\in\Delta_B}$ be the corresponding fundamental weights. To an element $\alpha \in \Delta_B$, we have a unique maximal parabolic subgroup P_α of G containing B such that $\Phi(N_{P_\alpha},T)\cap\Delta_B = \alpha$, where N_{P_α} is the unipotent radical of P_α. This gives a bijective correspondence between the simple roots in Δ_B and the maximal parabolic subgroups of G containing B. Any maximal parabolic subgroup P of G is conjugate to certain P_α by an element $w \in W$, the element $w\cdot\varpi_\alpha$ doesn't depend on the choice of w, we denote it by ϖ_P.

We use the (G,M) notation of Arthur. Let $\mathcal{F}(T)$ be the set of parabolic subgroups of G containing T, let $\mathcal{L}(T)$ be the set of Levi subgroups of G containing T. For every $M \in \mathcal{L}(T)$, we denote by $\mathcal{P}(M)$ the set of parabolic subgroups of G whose Levi factor is M. For $P \in \mathcal{P}(M)$, we denote by P^- the opposite of P with respect to M. Let $X^*(M) = \mathrm{Hom}(M,\mathbb{G}_m)$ and $\mathfrak{a}_M^* = X^*(M)\otimes\mathbf{R}$. The restriction $X^*(M)\to X^*(T)$ induces an injection $\mathfrak{a}_M^* \hookrightarrow \mathfrak{a}_T^*$. Let $(\mathfrak{a}_T^M)^*$ be the subspace of $\mathfrak{a}_T^*$ generated by $\Phi(M,T)$. We have the decomposition in direct sums

$$\mathfrak{a}_T^* = (\mathfrak{a}_T^M)^* \oplus \mathfrak{a}_M^*.$$

The canonical pairing

$$X_*(T)\times X^*(T)\to\mathbf{Z}$$

can be extended linearly to $\mathfrak{a}_T\times\mathfrak{a}_T^*\to\mathbf{R}$, with $\mathfrak{a}_T = X_*(T)\otimes\mathbf{R}$. For $M\in\mathcal{L}(T)$, let $\mathfrak{a}_T^M\subset\mathfrak{a}_T$ be the subspace orthogonal to $\mathfrak{a}_M^*$, and $\mathfrak{a}_M\subset\mathfrak{a}_T$ be the subspace orthogonal to $(\mathfrak{a}_T^M)^*$, then we have the decomposition

$$\mathfrak{a}_T = \mathfrak{a}_M\oplus\mathfrak{a}_T^M,$$

let π_M, π^M be the projections to the two factors.

We identify $X_*(T)$ with $T(F)/T(\mathcal{O})$ by sending χ to $\chi(\epsilon)$. With this identification, the canonical surjection $T(F)\to T(F)/T(\mathcal{O})$ can be viewed as

$$T(F)\to X_*(T). \tag{1}$$

We use Λ_G to denote the quotient of $X_*(T)$ by the coroot lattice of G (the subgroup of $X_*(T)$ generated by the coroots of T in G). We have a canonical homomorphism

$$G(F)\to\Lambda_G, \tag{2}$$

which is characterized by the following properties: it is trivial on the image of $G_{\mathrm{sc}}(F)$ in $G(F)$ (G_{sc} is the simply connected cover of the derived group of G), and its restriction to $T(F)$ coincides with the composition of (1) with the projection of $X_*(T)$ to Λ_G. Since the morphism (2) is trivial on $G(\mathcal{O})$, it descends to a map

$$\nu_G : \mathscr{X}\to\Lambda_G,$$

whose fibers are the connected components of $\mathscr{X}$.

Finally, we suppose that $\gamma\in\mathfrak{t}(\mathcal{O})$ satisfies $\gamma\equiv 0 \mod \epsilon$ to avoid unnecessary complications.

Acknowledgement. I want to thank Gérard Laumon for his encouragements during the preparation of this work.

2. The fundamental domain

2.1. **Truncated affine Springer fibers.** For $M \in \mathcal{L}(T)$, the natural inclusion of $M(F)$ in $G(F)$ induces a closed immersion of $\mathscr{X}^M$ in $\mathscr{X}^G$. For $P = MN \in \mathcal{F}(T)$, we have the retraction

$$f_P : \mathscr{X} \to \mathscr{X}^M$$

which sends $gK = nmK$ to $mM(\mathcal{O})$, where $g = nmk$, $n \in N(F)$, $m \in M(F)$, $k \in K$ is the Iwasawa decomposition. More generally we can define $f_{P_L}^L : \mathscr{X}^L \to \mathscr{X}^M$ for $L \in \mathcal{L}(T)$, $L \supset M$ and $P_L \in \mathcal{P}^L(M)$. These retractions satisfy the transition property: Suppose that $Q \in \mathcal{P}(L)$ satisfy $Q \supset P$, then

$$f_P = f_{P\cap L}^L \circ f_Q.$$

For $P \in \mathcal{F}(T)$, we have the function $H_P : \mathscr{X} \to \mathfrak{a}_M^G = \mathfrak{a}_M/\mathfrak{a}_G$ which is the composition

$$H_P : \mathscr{X} \xrightarrow{f_P} \mathscr{X}^M \xrightarrow{\nu_M} \Lambda_M \to \mathfrak{a}_M^G.$$

Proposition 2.1 (Arthur). *Let $B', B'' \in \mathcal{P}(T)$ be two adjacent Borel subgroups, let $\alpha_{B',B''}^\vee$ be the coroot which is positive with respect to B' and negative with respect to B''. Then for any $x \in \mathscr{X}$, we have*

$$H_{B'}(x) - H_{B''}(x) = n(x, B', B'') \cdot \alpha_{B',B''}^\vee,$$

with $n(x, B', B'') \in \mathbf{Z}_{\geq 0}$.

Proof. Let P be the parabolic subgroup generated by B' and B'', let $P = MN$ be the Levi factorization. The application $H_{B'}$ factor through f_P, i.e. we have commutative diagram

$$\begin{tikzcd}
\mathscr{X} \arrow[d, "f_P"'] \arrow[dr, "H_{B'}"] & \\
\mathscr{X}^M \arrow[r, "H_{B'\cap M}^M"'] & \mathfrak{a}_T^G
\end{tikzcd}$$

and similarly for $H_{B''}$. Since M has semisimple rank 1, the proposition is thus reduced to $G = \mathrm{SL}_2$. In this case, let T be the maximal torus of the diagonal matrices, $B' = \begin{pmatrix} * & * \\ & * \end{pmatrix}$, $B'' = \begin{pmatrix} * & \\ * & * \end{pmatrix}$, and we identify $\mathfrak{a}_T^G$ with the line $H = \{(x, -x) \mid x \in \mathbf{R}\} \subset \mathbf{R}^2$ in the usual way. By the Iwasawa decomposition, any point $x \in \mathscr{X}$ can be written as $x = \begin{pmatrix} a & b \\ & d \end{pmatrix} K$. Let $m = \min\{\mathrm{val}(a), \mathrm{val}(b)\}$, $n = \mathrm{val}(d)$, then $m + n \leq \mathrm{val}(a) + \mathrm{val}(d) = 0$ and

$$H_{B'}(x) = (-n, n), \quad H_{B''}(x) = (m, -m).$$

So

$$H_{B'}(x) - H_{B''}(x) = (-(n+m), n+m) = -(n+m)\cdot \alpha^{\vee}_{B',B''},$$

and the proposition follows. □

For any point $x \in \mathscr{X}$, we denote by $\mathrm{Ec}(x)$ the convex envelope in $\mathfrak{a}_T^G$ of the $H_{B'}(x)$, $B' \in \mathcal{P}(T)$. For any $P \in \mathcal{F}(T)$, we denote by $\mathrm{Ec}^P(x)$ the face of $\mathrm{Ec}(x)$ whose vertices are $H_{B'}(x)$, $B' \in \mathcal{P}(T)$, $B' \subset P$.

Definition 2.1. A family $D = (\lambda_{B'})_{B' \in \mathcal{P}(T)}$ of elements in $\mathfrak{a}_T^G$ is called (G,T)-*orthogonal* if it satisfies

$$\lambda_{B'} - \lambda_{B''} \in \mathbf{R}_{\geq 0}\cdot \alpha^{\vee}_{B',B''},$$

for any two adjacent Borel subgroups B', $B'' \in \mathcal{P}(T)$.

Let $D = (\lambda_{B'})_{B' \in \mathcal{P}(T)}$ be a (G,T)-orthogonal family, we also denote by D the convex envelope of $\lambda_{B'}$, $B' \in \mathcal{P}(T)$ in $\mathfrak{a}_T^G$. For $P = MN \in \mathcal{F}(T)$, parallel to $\mathrm{Ec}^P(x)$, we denote by D^P the face of D whose vertices are $\lambda_{B'}$, $B' \in \mathcal{P}(T)$, $B' \subset P$. With the projection π^M, it will also be seen as a (M,T)-orthogonal family.

Following Chaudouard and Laumon [CL], we define the *truncated affine grassmannian* $\mathscr{X}(D)$ to be

$$\mathscr{X}(D) = \{x \in \mathscr{X} \mid \mathrm{Ec}(x) \subset D\},$$

and the *truncated affine Springer fiber* $\mathscr{X}_\gamma(D)$ to be the intersection $\mathscr{X}_\gamma \cap \mathscr{X}(D)$. The connected components of $\mathscr{X}(D)$ and $\mathscr{X}_\gamma(D)$ are projective k-schemes. It should be noted that there is slight difference between different components of both $\mathscr{X}(D)$ and $\mathscr{X}_\gamma(D)$.

2.2. **The fundamental domain.** A point $x = gK \in \mathscr{X}_\gamma$ is said to be *regular* if the image of $\mathrm{Ad}(g^{-1})\gamma$ under the reduction $\mathfrak{g}(\mathcal{O}) \to \mathfrak{g}(k)$ is regular. It can be proved that the subvariety $\mathscr{X}_\gamma^{\mathrm{reg}}$ of regular points is open dense in $\mathscr{X}_\gamma$.

Proposition 2.2 (Bezrukavnikov). *The group $T(F)$ acts transitively on $\mathscr{X}_\gamma^{\mathrm{reg}}$.*

The reader is referred to [B] for the proof. As a consequence, the abelian group Λ generated by $\chi(\epsilon) \in T(F)$, $\chi \in X_*(T)$ acts freely and transitively on the irreducible components of $\mathscr{X}_\gamma$.

Goresky, Kottwitz and Macpherson [GKM3] have given the following description of regular points in $\mathscr{X}_\gamma$. We reproduce their proof here.

Lemma 2.3 (Goresky-Kottwitz-Macpherson). *A point $x \in \mathscr{X}_\gamma$ is regular if and only if for any Levi subgroup $M \in \mathcal{L}(T)$ of semisimple rank 1, the point $f_P(x) \in \mathscr{X}_\gamma^M$ is regular for any $P \in \mathcal{P}(M)$.*

Proof. For $x = gK \in \mathscr{X}$, the image of $\mathrm{Ad}(g)^{-1}\gamma$ under the reduction $\mathfrak{g}(\mathcal{O}) \to \mathfrak{g}(k)$ is well defined up to conjugacy, we denote it by $u_G(x)$. For any $P = MN \in \mathcal{F}(T)$, let $g = pk$, $p \in P(F)$, $k \in K$, then $\mathrm{Ad}(p)^{-1}\gamma \in \mathfrak{p}(F) \cap \mathfrak{g}(\mathcal{O}) = \mathfrak{p}(\mathcal{O})$. Its image in $\mathfrak{p}(k)$ under the reduction is well defined up to conjugacy, we will denote it by $u_P(x)$. It is obvious that $u_P(x)$ goes to $u_M(f_P(x))$ under the projection $\mathfrak{p} \to \mathfrak{m}$. So if $u_G(x)$ is regular, then

$u_M(f_P(x))$ is regular since $u_P(x)$ lies in the same conjugacy class as $u_G(x)$ in $\mathfrak{g}(k)$. This proves the necessary part of the lemma.

For the sufficiency, it is enough to prove that $u_B(x)$ is regular. For $\alpha \in \Delta_B$, let Q_α be the parabolic subgroup generated by B and $s_\alpha \cdot B$, where $s_\alpha \in W$ is the simple reflection associated to α. Let $Q_\alpha = M_\alpha N_\alpha$ be the Levi decomposition, then M_α is of semisimple rank 1. Now $u_B(x)$ goes to $u_{M_\alpha}(f_{Q_\alpha}(x))$ under the composition $\mathfrak{b} \hookrightarrow \mathfrak{q}_\alpha \twoheadrightarrow \mathfrak{m}_\alpha$. Since $u_{M_\alpha}(f_{Q_\alpha}(x))$ is regular in $\mathfrak{m}_\alpha$ for any $\alpha \in \Delta_B$, this implies that $u_B(x)$ is regular.

□

Proposition 2.4 (Goresky-Kottwitz-Macpherson). *Let $x \in \mathscr{X}_\gamma$.*

1. *For any two adjacent Borel subgroups $B', B'' \in \mathcal{P}(T)$, we have*
$$n(x, B', B'') \leq \mathrm{val}(\alpha_{B',B''}(\gamma)),$$
where $\alpha_{B',B''}$ is the root associated with the coroot $\alpha^\vee_{B',B''}$.
2. *The point x is regular if and only if the above equality holds for any two adjacent Borel subgroups.*

Proof. First of all, observe that for any $x, y \in \mathscr{X}$ such that y lies in the closure of the orbit $T(\mathcal{O}) \cdot x$, we have $\mathrm{Ec}(y) \subset \mathrm{Ec}(x)$. Now that $\mathscr{X}_\gamma^{\mathrm{reg}}$ is dense open in $\mathscr{X}_\gamma$, it suffices to prove the second assertion. By lemma 2.3, it suffices to prove the proposition for $G = \mathrm{GL}_2$. This follows from proposition 2.5, where we have picked a particular regular point $x_0 \in \mathscr{X}_\gamma^{\mathrm{reg}}$ and calculated that

$$H_B(x_0) = (\mathrm{val}(\alpha(\gamma)),\, 0), \quad H_{B^-}(x_0) = (0,\, \mathrm{val}(\alpha(\gamma))).$$

It is obvious that $H_B(x_0) - H_{B^-}(x_0) = \mathrm{val}(\alpha(\gamma)) \cdot \alpha^\vee$.

□

The above results motivate the following definition.

Definition 2.2. Take a regular point $x \in \mathscr{X}_\gamma^{\mathrm{reg}}$. Let F_γ be the connected component of the truncated affine Springer fiber $\mathscr{X}_\gamma\big((H_{B'}(x))_{B' \in \mathcal{P}(T)}\big)$ which contains x, it is *the fundamental domain* of $\mathscr{X}_\gamma$ with respect to the action of Λ.

It is clear that different choice of $x \in \mathscr{X}_\gamma^{\mathrm{reg}}$ gives rise to isomorphic fundamental domain.

2.3. **Examples for GL_d.** For the group GL_d, $\gamma \in \mathfrak{t}(\mathcal{O})$ regular, we have a particular choice of a regular point x_0 on $\mathscr{X}_\gamma$. Let $x_0 \in \mathscr{X}_\gamma^{\mathrm{GL}_d}$ be the point representing the lattice $\mathcal{O}[\gamma]$ sitting inside $F[\gamma] \cong F[X]/(X-\gamma_1) \oplus \cdots \oplus F[X]/(X-\gamma_d) \cong F^d$, where γ_i are the eigenvalues of γ. Taking $\{1, \gamma, \cdots, \gamma^{d-1}\}$ as a basis of $F[\gamma]$, we check easily that x_0 is a regular point.

Proposition 2.5. *For $\sigma \in \mathfrak{S}_d$, we have*

$$H_{\sigma B^-}(x_0) = \sigma^{-1}\left[\bigoplus_{i=1}^{d} \mathfrak{p}^{\sum_{j=1}^{i-1} \mathrm{val}(\alpha_{\sigma(i),\sigma(j)}(\gamma))}\right].$$

Proof. Let $\{e_1, \cdots, e_d\}$ be the natural basis of F^d, the vector $\gamma^s \in \mathcal{O}[\gamma]$ corresponds to the vector $\sum_{i=1}^d \gamma_i^s e_i$ in F^d. Let g be the matrix

$$\begin{bmatrix} 1 & 1 & \cdots & 1 \\ \gamma_1 & \gamma_2 & \cdots & \gamma_d \\ \vdots & \vdots & & \vdots \\ \gamma_1^{d-1} & \gamma_2^{d-1} & \cdots & \gamma_d^{d-1} \end{bmatrix},$$

then $\mathcal{O}[\gamma] = g\mathcal{O}^d$. From this expression, we see that it suffices to prove the proposition for the standard B^-.

After certain elementary operations on the columns, the matrix g can be put in lower triangular form with 1, $\gamma_2 - \gamma_1$, $(\gamma_3 - \gamma_2)(\gamma_3 - \gamma_1), \cdots, \prod_{i=1}^{d-1}(\gamma_d - \gamma_i)$ on the diagonal from top to bottom, from which the claim for $H_{B^-}(x_0)$ follows easily. □

Let $\gamma \in \mathfrak{t}(\mathcal{O})$ be regular in minimal form, suppose that its valuation data $(n_1, \cdots, n_{d-1})$ satisfies $n_1 \leq n_2 \leq \cdots \leq n_{d-1}$, then the fundamental domain F_γ can be written as the intersection of $\mathscr{X}_\gamma$ with two affine Schubert varieties. First of all, we identify $X_*(T)$ with $\mathbf{Z}^d$ in the natural way. For $\mu \in X_*(T)$, we denote by $\mathrm{Sch}(\mu)$ the affine Schubert variety $\overline{K\epsilon^\mu K/K}$. We fix

$$\begin{aligned} \mu &= \left(0,\, n_1,\, n_1 + n_2,\, \cdots,\, \sum_{i=1}^{d-1} n_i\right), \\ \lambda &= \left((d-1)n_1,\, n_1 + (d-2)n_2,\, n_1 + n_2 + (d-3)n_3,\, \cdots,\, \sum_{i=1}^{d-1} n_i,\, \sum_{i=1}^{d-1} n_i\right). \end{aligned}$$

Observe that

$$\mu_i = \sum_{j=1}^{i-1} \mathrm{val}(\alpha_{j,i}(\gamma)), \quad \lambda_i = \sum_{\substack{j=1 \\ j \neq i}}^{d} \mathrm{val}(\alpha_{j,i}(\gamma)). \tag{3}$$

Proposition 2.6. *In the above setting, the fundamental domain F_γ is the intersection*

$$F_\gamma = \mathscr{X}_\gamma \cap \left[\mathrm{Sch}(\mu) \cap \epsilon^\lambda \cdot \mathrm{Sch}(-\mu)\right].$$

Proof. Let D_1, D_2 be the convex envelope of the (G,T)-orthogonal family $(w \cdot \mu)_{w \in W}$ and $(\lambda - w \cdot \mu)_{w \in W}$ respectively. By Bruhat-Tits decomposition, we have

$$\mathscr{X}^{|\mu|}(D_1) = \mathrm{Sch}(\mu), \quad \mathscr{X}^{|\mu|}(D_2) = \epsilon^\lambda \cdot \mathrm{Sch}(-\mu),$$

where $\mathscr{X}^{|\mu|}$ is the connected component containing ϵ^μ. So we only need to prove that $\mathrm{Ec}(x_0) = D_1 \cap D_2$. Using proposition 2.5, on verifies that $H_{B'}(x_0) \in D_1 \cap D_2$, $\forall\, B' \in \mathcal{P}(T)$, so $\mathrm{Ec}(x_0) \subset D_1 \cap D_2$. Given a (G,T)-orthogonal family D, given $P = MN \in \mathcal{F}(T)$ maximal, let $d_P(D)$ be the distance between the two opposite faces D^P and D^{P^-}. It suffices then to prove that

$$d_P(\mathrm{Ec}(x_0)) = d_P(D_1 \cap D_2).$$

Choose a minimal gallery of Borel subgroups $B_1,\cdots,B_{l+1}$ such that $B_1\in P$, $B_{l+1}\in P^-$, then $\alpha_{B_i,B_{i+1}}$, $i=1,\cdots,l$ runs through $\Phi(N,T)$ exactly once. So we have

$$\begin{aligned} d_P(\mathrm{Ec}(x_0)) &= \varpi_P(H_{B_1}(x_0)-H_{B_{l+1}}(x_0))=\sum_{i=1}^{l}\varpi_P(H_{B_i}(x_0)-H_{B_{i+1}}(x_0)) \\ &= \sum_{\alpha\in\Phi(N,T)}\varpi_P(\mathrm{val}(\alpha(\gamma))\cdot\alpha^\vee)=\sum_{\alpha\in\Phi(N,T)}\mathrm{val}(\alpha(\gamma)). \end{aligned}$$

To calculate $d_P(D_1\cap D_2)$, let a_P be the distance of $D_1^{P^-}$ and D_2^P, then

$$d_P(D_1\cap D_2)=\min\{d_P(D_1),\,a_P\}.$$

Let $P_i=M_iN_i\in\mathcal{F}(T)$ be the maximal parabolic subgroup associated to the simple root α_i. For P conjugate to P_i or P_{d-i}, $i\le d/2$, we calculate that

$$d_P(D_1)=\sum_{j=1}^{i}\sum_{l=1+j}^{d-i+j-1}\mathrm{val}(\alpha_{l,d-i+j}(\gamma))=\sum_{j=1}^{i}\sum_{l=1+j}^{d-i+j-1}n_l.$$

We calculate a_{P_i} to be

$$\begin{aligned} a_{P_i} &= 2\varpi_{P_i^-}(\mu)-\varpi_{P_i^-}(\lambda)=-2\varpi_i(\mu)+\varpi_i(\lambda) \\ &= 2\left(\mu_{i+1}+\cdots+\mu_d-\frac{d-i}{d}(\mu_1+\cdots+\mu_d)\right) \\ &\quad -\left(\lambda_{i+1}+\cdots+\lambda_d-\frac{d-i}{d}(\lambda_1+\cdots+\lambda_d)\right) \\ &= \sum_{\alpha\in\Phi(N_i,T)}\mathrm{val}(\alpha(\gamma)), \end{aligned}$$

here we use equation (3) in the last equality. Conjugate the above calculation by $\sigma\in\mathfrak{S}_d$, we found that

$$a_P=\sum_{\alpha\in\Phi(N,T)}\mathrm{val}(\alpha(\gamma)).$$

It is easy to verify that $a_P\le d_P(D_1)$, so we have $d_P(D_1\cap D_2)=a_P=d_P(\mathrm{Ec}(x_0))$.

$\square$

3. ARTHUR-KOTTWITZ REDUCTION

Fix a regular point $x_0\in\mathscr{X}_\gamma^{\mathrm{reg}}$. Take $\xi\in\mathfrak{a}_T^G$ such that $\alpha(\xi)$ is positive but almost equal to 0 for any $\alpha\in\Delta_B$. Let $D_0=(\lambda_{B'})_{B'\in\mathcal{P}(T)}$ be the (G,T)-orthogonal family given by

$$\lambda_{B'}=H_{B'}(x_0)+w'\cdot\xi,$$

where $w' \in W$ is taken such that $B' = w' \cdot B$. For $P = MN \in \mathcal{F}(T)$, define R_P to be the subset of $\mathfrak{a}_T^G$ satisfying conditions

$$\begin{aligned} \pi^M(a) &\subset D_0^P; \\ \alpha(\pi_M(a)) &\geq \alpha(\pi_M(\lambda_{B'})), \forall\, \alpha \in \Phi(N,T), \forall\, B' \in \mathcal{P}(T),\ B' \subset P. \end{aligned}$$

Notice that $R_G = D_0$. We get a partition

$$\mathfrak{a}_T^G = \bigcup_{P \in \mathcal{F}(T)} R_P. \tag{4}$$

The figure 1 gives an illustration of the partition for the group GL_3. The partition (4) induces a disjoint partition of $\mathscr{X}_\gamma^T$ via the map $\mathscr{X}_\gamma^T \cong X_*(T) \to \mathfrak{a}_T^G$, since we have perturbed the (G,T)-family $(H_{B'}(x_0))_{B' \in \mathcal{P}(T)}$ with ξ.



Figure 1. Partition of $\mathfrak{a}_T^G$ for GL_3.

Lemma 3.1. *For any $x \in \mathscr{X}_\gamma$, there exists a unique $P \in \mathcal{F}(T)$ such that $\mathrm{Ec}^P(x) \subset R_P$.*

Proof. By proposition 2.4, the convex polytope $\mathrm{Ec}(x)$ is contained in a translation of $\mathrm{Ec}(x_0)$ by some $\lambda \in \Lambda$. The uniqueness is clear for such a translation, from which the uniqueness for general case follows.

Now we prove the existence. For a maximal parabolic $P \in \mathcal{F}(T)$, let $\overline{R}_P = \bigcup_{P' \subset P} R_P$. Notice that $\varpi_P(H_{B'}(x))$ doesn't depend on the choice of $B' \in \mathcal{P}(T)$, $B' \subset P$. Let P_0 be the maximal parabolic such that $\varpi_P(H_{B'}(x) - H_{B'}(x_0))$ is maximal among all the maximal parabolic subgroups P. It follows that $\mathrm{Ec}^{P_0}(x) \subset \overline{R}_{P_0}$, the basic reason is that in a right triangle with sides a, b, c, we always have $c > a, b$. Now we can use the retraction $f_{P_0} : \mathscr{X}_\gamma \to \mathscr{X}_\gamma^{M_{P_0}}$ to find the required parabolic subgroup P inductively.

□

With this lemma, we define $S_P := \{x \in \mathscr{X}_\gamma \mid \mathrm{Ec}^P(x) \subset R_P\}$. Notice that F_γ is one connected component of $S_G = \mathscr{X}_\gamma(D_0)$, the other connected components of S_G have slight difference from F_γ. We get a disjoint partition

$$\mathscr{X}_\gamma = \mathscr{X}_\gamma(D_0) \cup \bigcup_{\substack{P \in \mathcal{F}(T) \\ P \neq G}} S_P.$$

Consider the restriction of the retraction $f_P : \mathscr{X} \to \mathscr{X}^M$ to S_P, its image is $S_P \cap \mathscr{X}^M$. Recall that the connected components of $\mathscr{X}^M$ are fibers of the map $\nu_M : \mathscr{X}^M \to \Lambda_M$. For $\nu \in \Lambda_M$, let $\mathscr{X}^{M,\nu}$ be its fiber at ν. Let $S_P^\nu = f_P^{-1}(S_P \cap \mathscr{X}^{M,\nu})$, it is easy to verify that

$$S_P \cap \mathscr{X}^{M,\nu} = \mathscr{X}_\gamma^{M,\nu}(D_0^P).$$

Proposition 3.2. *The retraction $f_P : S_P^\nu \to \mathscr{X}_\gamma^{M,\nu}(D_0^P)$ is an iterated fibration in affine spaces.*

Proof. We follow the proof of Kazhdan-Lusztig [KL], §5. By assumption, $\mathrm{char}(k) > \mathrm{rk}(G)$, the exponential map $\exp : \mathfrak{n} \to N$ is well defined. The group N has the decreasing filtration by normal subgroups

$$N_0 = N \supset N_1 = [N, N] \supset \cdots \supset N_i = [N_{i-1}, N] \supset \cdots \supset N_{\mathrm{rk}(G)} \supset 1.$$

The exponential map induces an isomorphism $\mathfrak{n}_i/\mathfrak{n}_{i+1} \to N_i/N_{i+1}$ which sends n_i to $1+n_i$.

Since $f_P(ux) = f_P(x)$, $\forall\, u \in N$, $x \in \mathscr{X}$, by the definition of S_P^ν, we have the identity

$$S_P^\nu = [N(F) \cdot \mathscr{X}_\gamma^{M,\nu}(D_0^P)] \cap \mathscr{X}_\gamma.$$

So the fiber of $f_P : S_P^\nu \to \mathscr{X}_\gamma^{M,\nu}(D_0^P)$ at $mM(\mathcal{O})$ is

$$\Big\{umM(\mathcal{O}) \mid u \in N(F),\ \mathrm{Ad}(u^{-1})\gamma \in \mathrm{Ad}(m)\mathfrak{g}(\mathcal{O})\Big\}.$$

We'll prove that they form a family which is an iterated fibration in affine spaces.

Let $\mathcal{K}$ be the K-equivariant fiber bundle $G(F) \times_K K$ on $\mathscr{X}$, let $\mathfrak{K}$ be the K-equivariant vector bundle $G(F) \times_K \mathfrak{g}(\mathcal{O})$ on $\mathscr{X}$, where K acts on K and $\mathfrak{g}(\mathcal{O})$ by conjugaison. Let $\widetilde{N}_i$ be the constant fiber bundle $\mathscr{X} \times N_i(F)$, let $\tilde{\mathfrak{n}}_i$ be the constant vector bundle $\mathscr{X} \times \mathfrak{n}_i(F)$. We denote also $\widetilde{N} = \widetilde{N}_0$.

To begin with, observe that with the retraction f_P, the locally closed subvariety $f_P^{-1}(\mathscr{X}^{M,\nu})$ of $\mathscr{X}$ is isomorphic to the restriction of the fiber bundle $\widetilde{N}/\widetilde{N} \cap \mathcal{K}$ over $\mathscr{X}^{M,\nu}$, we will identify them in the following. For $i = 0, \cdots, \mathrm{rk}(G)+1$, let S_i be the sub bundle of $\widetilde{N}_i \backslash \widetilde{N}/\widetilde{N} \cap \mathcal{K}$ restricted to $\mathscr{X}_\gamma^{M,\nu}(D_0^P)$, whose fiber at $mM(\mathcal{O})$ is given by

$$\Big\{u \in N_i(F) \backslash N(F)/N(F) \cap \mathrm{Ad}(m)K \mid \mathrm{Ad}(u)^{-1}\gamma \in \mathrm{Ad}(m)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_i(F)\Big\}.$$

Let $p_i : S_{i+1} \to S_i$ be the natural projection, we get a tower of projections

$$S_P^\nu \cong S_{\mathrm{rk}(G)+1} \to S_{\mathrm{rk}(G)} \to \cdots \to S_0 \cong \mathscr{X}_\gamma^{M,\nu}(D_0^P).$$

The last isomorphism is due to the equivalence of the equations $\gamma \in \mathrm{Ad}(m)\mathfrak{g}(\mathcal{O})+\mathfrak{n}(F)$ and $\gamma \in \mathrm{Ad}(m)\mathfrak{g}(\mathcal{O})$ since $\mathrm{Ad}(m)^{-1}\gamma \in \mathfrak{m}(F)$. We will prove that each S_{i+1} is a homogeneous space under a vector bundle over S_i, this will end the proof of the proposition.

Given $gK \in S_i$, we have

$$\gamma \in \mathrm{Ad}(g)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_i(F).$$

Let $u = 1 + n \in N_{i+1}(F)\backslash N_i(F)$, with $n \in \mathfrak{n}_{i+1}(F)\backslash \mathfrak{n}_i(F)$, then

$$\begin{aligned} ugK \in S_{i+1} &\iff \mathrm{Ad}(u^{-1})\gamma \in \mathrm{Ad}(g)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_{i+1}(F) \\ &\iff \gamma + [\gamma, n] \in \mathrm{Ad}(g)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_{i+1}(F). \end{aligned} \tag{5}$$

Using the isomorphism

$$\frac{\mathrm{Ad}(g)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_i(F)}{\mathrm{Ad}(g)\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_{i+1}(F)} \cong \frac{\mathfrak{n}_i(F)/\mathfrak{n}_i(F) \cap \mathrm{Ad}(g)\mathfrak{g}(\mathcal{O})}{\mathfrak{n}_{i+1}(F)/\mathfrak{n}_{i+1}(F) \cap \mathrm{Ad}(g)\mathfrak{g}(\mathcal{O})},$$

let $\bar{\gamma}$ be the image of γ under the isomorphism, then the equation (5) means that n should satisfy the equation $\mathrm{ad}(\gamma)n = -\bar{\gamma}$ in the above quotient. Consider the endomorphism $\mathrm{ad}(\gamma)$ of the restriction of the vector bundle

$$\frac{\tilde{\mathfrak{n}}_i/\tilde{\mathfrak{n}}_i \cap \mathfrak{K}}{\tilde{\mathfrak{n}}_{i+1}/\tilde{\mathfrak{n}}_{i+1} \cap \mathfrak{K}} \tag{6}$$

on S_i. It is surjective since $\mathrm{ad}(\gamma) : \mathfrak{n}_i(F) \to \mathfrak{n}_i(F)$ is. This means that there is always n such that equation (5) is satisfied, i.e. p_i are surjective for all i. Further more, let V_i be kernel of the endomorphism $\mathrm{ad}(\gamma)$ of the vector bundle (6), then V_i is a vector bundle on S_i. The above calculation shows that S_{i+1} is a homogeneous space over S_i under the vector bundle V_i.

□

Proposition 3.3. *The stratas S_P^ν are locally closed sub varieties of $\mathscr{X}_\gamma$. Further more, in the decomposition*

$$\mathscr{X}_\gamma = \mathscr{X}_\gamma(D_0) \cup \bigcup_{\substack{P \in \mathcal{F}(T) \\ P \neq G}} \bigcup_{\nu \in \Lambda_{M_P} \cap R_P} S_P^\nu, \tag{7}$$

we can order the strata S_P^ν such that at each step we have a closed sub variety of $\mathscr{X}_\gamma$.

Proof. To begin with, $\mathscr{X}_\gamma(D)$ is a closed sub variety of $\mathscr{X}_\gamma$ for any (G,T)-orthogonal family D. Now we prove by induction. Let $P_0 = M_0N_0 \in \mathcal{F}(T)$ be a maximal parabolic subgroup containing P. For $P' = M'N' \in \mathcal{F}(T)$, $P' \subset P_0$, let $p_{M_0}^{M'}$ be the natural projection $\Lambda_{M'} \to \Lambda_{M_0}$, let $\nu_0 = p_{M_0}^{M}(\nu)$. Consider

$$Z_{P_0}^{\nu_0} := \bigcup_{P' \subset P_0} \bigcup_{p_{M_0}^{M'}(\nu') = \nu_0} S_{P'}^{\nu'}.$$

Firstly, $Z_{P_0}^{\nu_0}$ can be written as a difference $\mathscr{X}_\gamma(D)\backslash\mathscr{X}_\gamma(D')$ for two (G,T)-orthogonal family D, D'. Secondly, observe that

$$S_{P'}^{\nu'} = [N_0(F) \cdot (S_{P'}^{\nu'} \cap \mathscr{X}^{M_0,\nu_0})] \cap \mathscr{X}_\gamma,$$

the same proof as that of proposition 3.2 shows that the retraction

$$f_{P_0} : Z_{P_0}^{\nu_0} \to \mathscr{X}_\gamma^{M_0,\nu_0}(D^{P_0})$$

is an iterated fibration in affine spaces. Now the claim follows by induction. □

By proposition 3.2, each strata S_P^ν has an iterated affine fibration onto $\mathscr{X}_\gamma^{M_P,\nu}(D_0^P)$, so the study of $\mathscr{X}_\gamma$ is reduced to that of F_γ. We call the decomposition (7) *the Arthur-Kottwitz reduction*. We remark that this is not a reduction of Harder-Narasimhan type.

Lemma 3.4. *Suppose that F_γ^M is cohomologically pure for any Levi subgroup M of G containing T. Suppose that F_γ is pure, then the truncated affine Springer fiber $\mathscr{X}_\gamma^\nu(D_0)$ is pure for all $\nu \in \Lambda_G$.*

Proof. After certain translation on $\mathscr{X}$ by Λ, we have $F_\gamma = \mathscr{X}_\gamma^\nu(D_0 + \varpi^\vee)$, where $\varpi^\vee$ is a minuscule coweight, and $D_0 + \varpi^\vee$ is the translation of D_0 by $\varpi^\vee$. It is easy to see that $\mathscr{X}_\gamma^\nu(D_0) \subset F_\gamma$. Applying the reduction of Arthur-Kottwitz, the open sub variety $F_\gamma \backslash \mathscr{X}_\gamma^\nu(D_0)$ is naturally stratified into finite unions of $S_P^{\nu'} \cap F_\gamma$. Since the two truncation parameters differ by a minuscule coweight, we have an inclusion of T-fixed points $(S_P^{\nu'})^T \subset (F_\gamma)^T$ and $(S_P^{\nu'})^T$ all lies on the faces of $\mathrm{Ec}(x_0)$. By proposition 2.4, we have $S_P^{\nu'} \subset F_\gamma$, so $S_P^{\nu'} \cap F_\gamma = S_P^{\nu'}$.

Now we prove by induction. Suppose that the lemma is proved for all the Levi subgroups $M \in \mathcal{L}(T)$, then $\mathscr{X}_\gamma^{M,\nu'}(D_0^P)$ are all cohomologically pure for all $P \in \mathcal{P}(M)$ and all $\nu' \in \Lambda_M$. By proposition 3.2 and 3.3, we see that $F_\gamma \backslash \mathscr{X}_\gamma^\nu(D_0)$ is cohomologically pure. Now the long exact sequence

$$\cdots \to H^{i-1}(\mathscr{X}_\gamma^\nu(D_0)) \to H_c^i(F_\gamma \backslash \mathscr{X}_\gamma^\nu(D_0)) \to H^i(F_\gamma) \to H^i(\mathscr{X}_\gamma^\nu(D_0)) \to \cdots$$

will split into short exact sequence

$$0 \to H_c^i(F_\gamma \backslash \mathscr{X}_\gamma^\nu(D_0)) \to H^i(F_\gamma) \to H^i(\mathscr{X}_\gamma^\nu(D_0)) \to 0,$$

because $H^{i-1}(\mathscr{X}_\gamma^\nu(D_0))$ is of weight at most $i-1$ by [Weil II]. The claim then follows from the above short exact sequence.

□

Now we come to the proof of theorem 1.1. We will prove a slightly stronger result. A (G,T)-orthogonal family $D = (\mu_{B'})_{B' \in \mathcal{P}(T)}$ is said to be *regular* with respect to D_0 if $\mu_{B'} \in R_{B'}$, $\forall\, B' \in \mathcal{P}(T)$.

Theorem 3.5. *Suppose that F_γ^M is cohomologically pure for any Levi subgroup M of G containing T. Let D be a (G,T)-orthogonal family regular with respect to D_0. Then F_γ is cohomologically pure if and only if the truncated affine Springer fiber $\mathscr{X}_\gamma(D)$ is.*

Proof. The complication that some connected components of $\mathscr{X}_\gamma(D)$ doesn't contain F_γ is already treated in lemma 3.4, so we can suppose that every connected component of $\mathscr{X}_\gamma(D)$ contains a translation of F_γ. Applying the Arthur-Kottwitz reduction to every connected component $\mathscr{X}_\gamma^\nu(D)$, we get a stratification of $\mathscr{X}_\gamma^\nu(D) \backslash F_\gamma$ into finite union of $S_P^{\nu'} \cap \mathscr{X}_\gamma^\nu(D)$.

The hypothesis that D is regular with respect to D_0 implies that each $S_P^{\nu'}$ is either contained in $\mathscr{X}_\gamma^\nu(D)$ or disjoint from it. Applying lemma 3.4 to the Levi subgroups $M \in \mathcal{L}(T)$, we see that all the truncated affine Springer fibers $\mathscr{X}_\gamma^{M,\nu'}(D_0^P)$, $P \in \mathcal{P}(M)$ are cohomologically pure, which implies that $\mathscr{X}_\gamma^\nu(D)\backslash F_\gamma$ is cohomologically pure by proposition 3.2 and 3.3. Now the theorem follows from the same argument as the last part of the proof of lemma 3.4.

□

4. RATIONALITY CONJECTURE FOR GL_3

Let $G = \mathrm{GL}_3$, let $\mathbf{n} = (n_1, n_2) \in \mathbf{N}^2$, $n_1 \leq n_2$, let $\gamma \in \mathfrak{t}(\mathcal{O})$ be in minimal form with root valuation $\mathbf{n}$. Let F_γ be the fundamental domain of $\mathscr{X}_\gamma$ containing x_0, where x_0 is the regular point defined in §2.3.

Proposition 4.1. *The fundamental domain F_γ is the intersection of $\mathscr{X}_\gamma$ with*

$$\mathrm{Sch}(2n_1 + n_2, 0, 0) \cap \mathrm{diag}(\epsilon^{-n_2}, \epsilon^{-n_1}, \epsilon^{-n_1}) \cdot \mathrm{Sch}(2n_1 + n_2, 2n_1 + n_2, 0).$$

Proof. By proposition 2.5, $\mathrm{Ec}(x_0)$ is the hexagon with vertices marked as indicated in figure 2.



FIGURE 2. Hexagon as intersection of two triangles.

This hexagon can also be represented as the intersection of two triangles as indicated also in the figure. Let △, ▽ be the upward and the downward triangle in the figure. By

Bruhat-Tits decomposition, we see that $\mathrm{Ec}(x) \in \nabla$ if and only if $x \in \mathrm{Sch}(2n_1+n_2,0,0)$. We notice that $\triangle$ is the translation by $(-n_2,-n_1,-n_1)$ of the triangle $\triangle'$ with vertices

$$(2n_1+n_2,2n_1+n_2,0), \quad (2n_1+n_2,0,2n_1+n_2), \quad (0,2n_1+n_2,2n_1+n_2).$$

Again by Bruhat-Tits decomposition, we see that $\mathrm{Ec}(x) \in\triangle'$ if and only if $x \in \mathrm{Sch}(2n_1+n_2,2n_1+n_2,0)$, the result follows directly from these considerations.

□

4.1. **Affine paving.** We can pave F_γ in affine spaces, the strategy is the following: By proposition 4.1, we have

$$F_\gamma = \mathscr{X}_\gamma \cap \mathrm{Sch}(2n_1+n_2,0,0) \cap \mathrm{diag}(\epsilon^{-n_2},\epsilon^{-n_1},\epsilon^{-n_1}) \cdot \mathrm{Sch}(2n_1+n_2,2n_1+n_2,0).$$

So we firstly pave the intersection of the two affine schubert varieties in affine spaces, but this paving doesn't induce an affine paving of F_γ, we need to regroup the resulting pavements and do a second nonstandard paving. Our main result in this section is:

Theorem 4.2. *The fundamental domain F_γ admits an affine paving, which only depends on the root valuation of γ.*

Proof. Let I be the standard Iwahori subgroup, i.e. it is the inverse image of the Borel subgroup B under the reduction $G(\mathcal{O}) \to G(k)$. Let $I' = \mathrm{Ad}(\mathrm{diag}(\epsilon^{n_1},\epsilon^{n_2},\epsilon^{n_2}))I$. By [C] corollary 2.3, we have the affine paving

$$\begin{aligned}
&\mathrm{Sch}(2n_1+n_2,0,0) \cap \mathrm{diag}(\epsilon^{-n_2},\epsilon^{-n_1},\epsilon^{-n_1}) \cdot \mathrm{Sch}(2n_1+n_2,2n_1+n_2,0)\\
&= \bigsqcup_{\mu\in(F_\gamma)^T} \mathrm{Sch}(2n_1+n_2,0,0) \cap I'\epsilon^\mu K/K\\
&= \bigsqcup_{\mu\in(F_\gamma)^T} \begin{bmatrix} \mathcal{O} & \mathfrak{p}^a & \mathfrak{p}^b \\ \mathfrak{p}^{n_2-n_1+1} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p}^{n_2-n_1+1} & \mathfrak{p} & \mathcal{O} \end{bmatrix} \epsilon^\mu K/K,
\end{aligned}$$

where $a=\max\{n_1-n_2,-\mu_2\}$, $b=\max\{n_1-n_2,-\mu_3\}$. We denote by $C(\mu)$ the resulting pavement containing ϵ^μ.

To pave F_γ, we cut it into 4 parts. Let $(\mu_1',\mu_2',\mu_3')=(\mu_1-n_1,\mu_2-n_2,\mu_3-n_2)$, and

$$\begin{aligned}
R_1 &= \{\mu \in (F_\gamma)^T \mid \mu_1' \leq \mu_2',\mu_3'\},\\
R_1' &= \{\mu \in (F_\gamma)^T \mid \mu_1' \geq \mu_2',\mu_3';\ \mu_2 \leq n_2-n_1;\ \mu_3 \leq n_2-n_1\},\\
R_2 &= \{\mu \in (F_\gamma)^T \mid \mu_2' < \mu_1',\mu_3';\ \mu_3 > n_2-n_1\},\\
R_3 &= \{\mu \in (F_\gamma)^T \mid \mu_3' < \mu_1',\mu_2';\ \mu_2 > n_2-n_1\}.
\end{aligned}$$

Although R_1 and R_1' may intersect at one point, it doesn't cause trouble to the paving. Figure 3 gives an idea of the cutting. Let $V_i=\bigsqcup_{\mu\in R_i} C(\mu)$, $i=1,2,3$. For $l\in\mathbf{Z}$, let $R_{i,l}=\{\mu\in R_i \mid \mu_i=l\}$ and $V_{i,l}=\bigsqcup_{\mu\in R_{i,l}} C(\mu)$. Similar notations for R_1'.



FIGURE 3. Nonstandard paving.

We use the Iwahori subgroup I' to pave $V_1 \cap \mathscr{X}_\gamma$. Since we have

$$C(\mu) = \begin{bmatrix} \mathcal{O} & & \\ \mathfrak{p}^{n_2-n_1+1} & \mathcal{O} & \mathcal{O} \\ \mathfrak{p}^{n_2-n_1+1} & \mathfrak{p} & \mathcal{O} \end{bmatrix} \epsilon^\mu K/K,$$

we see easily that $C(\mu) \cap \mathscr{X}_\gamma$ is isomorphic to an affine space.

We also use I' to pave $V_1' \cap \mathscr{X}_\gamma$. We have

$$C(\mu) = \begin{bmatrix} \mathcal{O} & \mathfrak{p}^{-\mu_2} & \mathfrak{p}^{-\mu_3} \\ & \mathcal{O} & \mathcal{O} \\ & \mathfrak{p} & \mathcal{O} \end{bmatrix} \epsilon^\mu K/K.$$

It is easily checked that $C(\mu) \cap \mathscr{X}_\gamma$ is isomorphic to an affine space.

We need a second nonstandard paving in order to pave $V_2 \cap \mathscr{X}_\gamma$ and $V_3 \cap \mathscr{X}_\gamma$. Since they are symmetric, we only give details for $V_3 \cap \mathscr{X}_\gamma$. Since $V_3 = \bigsqcup_{l \in \mathbf{Z}} V_{3,l}$, we only need to pave $V_{3,l} \cap \mathscr{X}_\gamma$. Let $I_l' = \mathrm{Ad}(\mathrm{diag}(1, \epsilon^l, \epsilon^l))I'$, we claim that

$$V_{3,l} \cap \mathscr{X}_\gamma = \bigsqcup_{\mu \in R_{3,l}} V_{3,l} \cap I_l' \epsilon^\mu K/K \cap \mathscr{X}_\gamma$$

is an affine paving. Since we have

$$C(\mu) = \begin{bmatrix} \mathcal{O} & \mathfrak{p}^{n_1-n_2} & \mathfrak{p}^{-\mu_3} \\ \mathfrak{p}^{n_2-n_1+1} & \mathcal{O} & \mathcal{O} \\ & \mathfrak{p} & \mathcal{O} \end{bmatrix} \epsilon^\mu K/K,$$

we see easily that $V_{3,l}$ admits an affine fibration onto the closed subvariety

$$\bigsqcup_{\mu\in R_{3,l}}\begin{bmatrix}\mathcal{O} & \mathfrak{p}^{n_1-n_2} & \\ \mathfrak{p}^{n_2-n_1+1} & \mathcal{O} & \\ & & \mathcal{O}\end{bmatrix}\epsilon^{\mu}K/K$$

of $\mathscr{X}^{\mathrm{GL}_2\times\mathrm{GL}_1}$. This implies

$$V_{3,l}\cap I'_l\epsilon^{\mu}K/K=\begin{bmatrix}\mathcal{O} & \mathfrak{p}^{c} & \mathfrak{p}^{-l}\\ \mathfrak{p}^{n_2-n_1+l+1} & \mathcal{O} & \mathcal{O}\\ & \mathfrak{p} & \mathcal{O}\end{bmatrix}\epsilon^{\mu}K/K,$$

with $c=\max(n_1-n_2-l,-\mu_2)$. With this equality, it is easily checked that $V_{3,l}\cap I'_l\epsilon^{\mu}K/K\cap\mathscr{X}_\gamma$ is isomorphic to an affine space.

It remains to precise the order of the paving. First of all, the Bruhat-Tits order with respect to I' induces an ordering of $V'_{1,l}$ and $V_{i,l}$, $i=1,2,3$, $l\in\mathbf{Z}$. On $V_{1,l}$ and $V'_{1,l}$ we use the Bruhat-Tits order with respect to I', while on $V_{2,l}$ and $V_{3,l}$, we use the Bruhat-Tits order with respect to I'_l.

□

4.2. **Rationality conjecture.** To calculate the Poincaré polynomial of F_γ, we proceed by an indirect way in order to avoid the combinatorial complexity. Our strategy is the following: we calculate firstly the Poincaré polynomial of $\mathscr{X}_\gamma\cap\mathrm{Sch}(2n_1+n_2,0,0)$, then we calculate the Poincaré polynomial of the complementary of F_γ, their difference gives what we want. It turns out that the complementary of F_γ can be paved in affine spaces.

Theorem 4.3. *The Poincaré polynomial of F_γ is*

$$P_{\mathbf{n}}(t)=\sum_{i=1}^{n_1}i(t^{4i-2}+t^{4i-4})+\sum_{i=2n_1}^{n_1+n_2-1}(2n_1+1)t^{2i}+\sum_{i=n_1+n_2}^{2n_1+n_2-1}4(2n_1+n_2-i)t^{2i}+t^{4n_1+2n_2}.$$

Taking into account the fact that F_{n_2,n_1} has the same Poincaré polynomial as F_{n_1,n_2}, we get the precise expression for the generating series.

Corollary 4.4. *The power series*

$$\sum_{n_1=1}^{+\infty}\sum_{n_2=1}^{+\infty}P_{(n_1,n_2)}(t)\,T_1^{n_1}T_2^{n_2}\in\mathbf{Z}[t][[T_1,T_2]]$$

equals the rational function

$$\begin{aligned}2\Bigg\{&\frac{(t^2+1)T_1T_2}{(1-T_2)(1-T_1T_2)(1-t^4T_1T_2)^2}+\frac{t^4T_1T_2^2(3-t^4T_1T_2)}{(1-T_2)(1-t^2T_2)(1-t^4T_1T_2)^2}\\ +\ &\frac{4t^4T_1T_2}{(1-t^2T_2)(1-t^4T_1T_2)^2(1-t^6T_1T_2)}+\frac{t^6T_1T_2}{(1-t^2T_2)(1-t^6T_1T_2)}\Bigg\}\\ -\ &\left[\frac{(t^2+1)T_1T_2}{(1-T_1T_2)(1-t^4T_1T_2)^2}+\frac{4t^4T_1T_2}{(1-t^4T_1T_2)^2(1-t^6T_1T_2)}+\frac{t^6T_1T_2}{1-t^6T_1T_2}\right]\end{aligned}$$

Proof of the theorem 4.3. To pave $\mathscr{X}_\gamma \cap \mathrm{Sch}(2n_1+n_2,0,0)$, we use the same idea as the proof of theorem 3.11 in [C]. We can pave $\mathrm{Sch}(2n_1+n_2,0,0)$ in affine spaces with the Iwahori subgroup

$$I' = \mathrm{Ad}(\mathrm{diag}(\epsilon^{2n_1+n_2},1,1))I.$$

Let $C(\mu)=\mathrm{Sch}(2n_1+n_2,0,0)\cap I'\epsilon^\mu K/K$, then we have the affine paving

$$\mathrm{Sch}(2n_1+n_2,0,0)=\bigsqcup_{\mu\in \mathrm{Sch}(2n_1+n_2,0,0)^T} C(\mu),$$

with

$$C(\mu)=\begin{bmatrix}\mathcal{O} & & \\ \mathfrak{p}^{-\mu_1} & \mathcal{O} & \mathcal{O}\\ \mathfrak{p}^{-\mu_1} & \mathfrak{p} & \mathcal{O}\end{bmatrix}\epsilon^\mu K/K.$$

Then we prove with the same method that $C(\mu)\cap\mathscr{X}_\gamma$ is an affine space of dimension

$$\min\{n_1,\mu_2\}+\min\{n_1,\mu_3\}+\min\left\{n_2,|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}\right\}.$$

It suffices to count the number of affine pavements of each dimension to get the Poincaré polynomial. To facilitate the work, we cut $\mathrm{Sch}(2n_1+n_2,0,0)^T$ into 7 parts, as indicated in figure 4, where

$$\begin{aligned}
R_1 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_2-\mu_3>n_2\},\\
R_1' &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3-\mu_2>n_2\},\\
R_2 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_2-\mu_3\le n_2, \mu_3<n_1, \mu_2>n_1\},\\
R_2' &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3-\mu_2\le n_2, \mu_2<n_1, \mu_3>n_1\},\\
R_3 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3\ge n_1, \mu_2\ge n_1\},\\
R_4 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3\le n_1, \mu_2\le n_1, n_2<\mu_1\le n_1+n_2\},\\
R_4' &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3<n_1, \mu_2<n_1, n_1+n_2<\mu_1\le 2n_1+n_2\},
\end{aligned}$$

Now we count the contribution of each part. We first sum over each blue lines as indicated in figure 4, then we add up all blue lines. We use the notation $\sum_{\mu=\nu}^{\nu'}$ to mean summation over the line having ends in ν, ν'. Since the Poincaré polynomial is a polynomial in t^2, we use $q:=t^2$ to simplify the notation.

(1) The contribution of $C(\mu)\cap\mathscr{X}_\gamma, \mu\in R_1$ to the Poincaré polynomial of $\mathscr{X}_\gamma\cap\mathrm{Sch}(2n_1+n_2,0,0)$ is

$$\sum_{i=0}^{n_1-1}\sum_{\mu=(0,2n_1+n_2-i,i)}^{(2n_1-2i-1,n_2+i+1,i)} q^{n_1+n_2+i}=2\sum_{i=1}^{n_1} iq^{2n_1+n_2-i}.$$

(2) The contribution of $C(\mu)\cap\mathscr{X}_\gamma, \mu\in R_1'$ is the same as R_1.



FIGURE 4. Partition of the triangle.

(3) The contribution of $C(\mu)\cap\mathscr{X}_\gamma, \mu\in R_2$ is

$$\begin{aligned}&\sum_{i=0}^{n_1-1}\sum_{\mu=(2n_1-2i,n_2+i,i)}^{(n_1+n_2-i-1,n_1+1,i)} q^{i+n_1+\mu_2-\mu_3}\\ =\ & n_1\sum_{i=2n_1+1}^{n_1+n_2} q^i+\sum_{i=1}^{n_1-1}(n_1-i)q^{n_1+n_2+i}.\end{aligned}$$

(4) The contribution of R'_2 is

$$\begin{aligned}&\sum_{i=0}^{n_1-1}\sum_{\mu=(2n_1-2i,i,n_2+i)}^{(n_1+n_2-i-1,i,n_1+1)} q^{i+n_1+\mu_3-\mu_2-1}\\ =\ & n_1\sum_{i=2n_1+1}^{n_1+n_2} q^{i-1}+\sum_{i=1}^{n_1-1}(n_1-i)q^{n_1+n_2+i-1}.\end{aligned}$$

(5) The contribution of R_3 is

$$\begin{aligned}
&\sum_{i=0}^{n_2}\sum_{\mu=(i,n_1,n_1+n_2-i)}^{(i,n_1+n_2-i,n_1)} q^{2n_1+|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}}\\
=\;& \sum_{i=0}^{n_2} q^{2n_1}(1+q+q^2+\cdots+q^{n_2-i})\\
=\;& q^{2n_1}\sum_{i=0}^{n_2}(n_2+1-i)q^i.
\end{aligned}$$

(6) The contribution of R_4 is

$$\begin{aligned}
&\sum_{i=0}^{n_1-1}\sum_{\mu=(n_1+n_2-i,n_1,i)}^{(n_1+n_2-i,i,n_1)} q^{n_1+i+|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}}\\
=\;& \sum_{i=0}^{n_1-1} q^{n_1+i}(1+q+\cdots+q^{n_1-i})\\
=\;& n_1q^{2n_1}+\sum_{i=0}^{n_1-1}(i+1)q^{n_1+i}.
\end{aligned}$$

(7) The contribution of R_4' is

$$\begin{aligned}
&\sum_{i=0}^{n_1-1}\sum_{\mu=(2n_1+n_2-i,0,i)}^{(2n_1+n_2-i,i,0)} q^{i+|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}}\\
=\;& \sum_{i=0}^{n_1-1} q^{i}(1+q+\cdots+q^{i}).
\end{aligned}$$

The complementary of F_γ in $\mathscr{X}_\gamma\cap\mathrm{Sch}(2n_1+n_2,0,0)$ can be paved in affine spaces in the following way: Observe that F_γ is contained in the intersection $\mathscr{X}_\gamma\cap\mathrm{Sch}(n_1+n_2,n_1,0)$, whose complementary in $\mathscr{X}_\gamma\cap\mathrm{Sch}(2n_1+n_2,0,0)$ can be paved in affine spaces using the standard Iwahori subgroup I. It suffices to pave the complementary of F_γ in $\mathscr{X}_\gamma\cap\mathrm{Sch}(n_1+n_2,n_1,0)$, which can be done by using the Iwahori subgroup

$$I''=\mathrm{Ad}(\mathrm{diag}(\epsilon^{n_1},\epsilon^{n_2},\epsilon^{n_2}))I.$$

We cut the complementary of F_γ^T in $\mathrm{Sch}(2n_1+n_2,0,0)^T$ as indicated in figure 5, where

$$\begin{aligned}
T_1 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_1\geq n_1+n_2+1\},\\
T_2 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_2\geq n_1+n_2+1\},\\
T_3 &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid \mu_3\geq n_1+n_2+1\},\\
T_1' &= \{\mu\in\mathrm{Sch}(2n_1+n_2,0,0)^T \mid 2n_1+1\leq \mu_1\leq n_1+n_2\}.
\end{aligned}$$



FIGURE 5. Complementary of F_γ.

The complementary of $\mathscr{X}_\gamma \cap \mathrm{Sch}(n_1+n_2,n_1,0)$ in $\mathscr{X}_\gamma \cap \mathrm{Sch}(2n_1+n_2,0,0)$ is

$$\bigsqcup_{\mu\in T_1\cup T_2\cup T_3} I\epsilon^\mu K/K\cap\mathscr{X}_\gamma.$$

It is easy to verify that this is an affine paving. To calculate its Poincaré polynomial, in each region we first sum over the vertices on the blue lines as indicated in figure 5, then we sum over all the lines.

(1) The contribution of T_1 is

$$\begin{aligned}&\sum_{i=0}^{n_1-1}\sum_{\mu=(2n_1+n_2-i,0,i)}^{(2n_1+n_2-i,i,0)} q^{2n_1+|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}}\\ =&\sum_{i=0}^{n_1-1} q^{2n_1}(1+q+\cdots+q^i).\end{aligned}$$

(2) The contribution of T_2 is

$$\sum_{i=0}^{n_1-1}\sum_{\mu=(0,2n_1+n_2-i,i)}^{(n_1-i-1,n_1+n_2+1,i)} q^{n_1+n_2+i}=\sum_{i=1}^{n_1} iq^{2n_1+n_2-i}.$$

(3) The contribution of T_3 is the same as that of T_2.

It remains to calculate the Poincaré polynomial of the complementary of F_γ in $\mathscr{X}_\gamma \cap \mathrm{Sch}(n_1+n_2, n_1, 0)$. By proposition 4.1, it is the union

$$\bigsqcup_{\mu\in T_1'} I''\epsilon^{\mu}K/K \cap \mathrm{Sch}(n_1+n_2, n_1, 0) \cap \mathscr{X}_\gamma.$$

By proposition 2.4, points in $I\epsilon^{\mu}K/K \cap \mathscr{X}_\gamma$, $\mu \in T_2 \cup T_3$ don't belong to any $B'\epsilon^{\nu}K/K \cap \mathscr{X}_\gamma$, $\nu \in T_1'$, for any $B' \in \mathcal{F}(T)$. The above intersection is thus equal to

$$\bigsqcup_{\mu\in T_1'} I''\epsilon^{\mu}K/K \cap \mathrm{Sch}(2n_1+n_2, 0, 0) \cap \mathscr{X}_\gamma,$$

which is easily verified to be an affine space of dimension

$$2n_1 + |\mu_2-\mu_3| + \frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2},$$

using the equality

$$I''\epsilon^{\mu}K/K \cap \mathrm{Sch}(2n_1+n_2,0,0) = \begin{bmatrix} \mathcal{O} & \mathfrak{p}^a & \mathfrak{p}^b \\ & \mathcal{O} & \mathcal{O} \\ & \mathfrak{p} & \mathcal{O} \end{bmatrix} \epsilon^{\mu}K/K,$$

where $a = \max\{n_1-n_2, -\mu_2\}$, $b = \max\{n_1-n_2, -\mu_3\}$.

Summing up the contributions of all the pavements in T_1' in the order as for the region T_1, we find the Poincaré polynomial of the complementary of F_γ in $\mathscr{X}_\gamma \cap \mathrm{Sch}(n_1+n_2, n_1, 0)$ to be

$$\begin{aligned} &\sum_{i=n_1}^{n_2-1} \sum_{\mu=(2n_1+n_2-i,0,i)}^{(2n_1+n_2-i,i,0)} q^{2n_1+|\mu_2-\mu_3|+\frac{\mathrm{sign}(\mu_2-\mu_3)-1}{2}} \\ = &\sum_{i=n_1}^{n_2-1} q^{2n_1}(1+q+\cdots+q^i). \end{aligned}$$

Now taking into account all the above calculations, we get the result as claimed in the theorem. ☐

Remark 4.1. Observe that in the above proof we actually give an affine paving of the complementary of F_γ in $\mathscr{X}_\gamma \cap \mathrm{Sch}(2n_1+n_2, 0, 0)$, and this paving can also be obtained by the Arthur-Kottwitz reduction. With more efforts, the same method can be generalized to GL_4 using pavings in [C].

REFERENCES

[A] J. Arthur, *The characters of discrete series as orbital integrals*, Invent. math. **32**(1976), 205-261.

[B] R. Bezrukavnikov, *The dimension of the fixed point set on affine flag manifolds*, Math. Res. Lett. 3 (1996), 185–189.

[C] Z. Chen, *Pureté des fibres de Springer affines pour* GL_4, `http://arxiv.org/pdf/1111.3352`, à paraitre dans Bull. SMF.

[CL] P-H. Chaudouard, G. Laumon, *Sur l'homologie des fibres de Springer affines tronquées*, Duke Math. J. 145 (2008), no. 3, 443–535.

[GKM1] M. Goresky, R. Kottwitz, R. Macpherson, *Homology of affine Springer fibers in the unramified case*, Duke Math. J. 121 (2004), no. 3, 509-561.

[GKM2] M. Goresky, R. Kottwitz, R. Macpherson, *Purity of equivalued affine Springer fibers*, Represent. Theory 10 (2006), 130-146.

[GKM3] M. Goresky, R. Kottwitz, R. Macpherson, *Regular points in affine Springer fibers*, Michigan Math. J. 53 (2005), no. 1, 97-107.

[GKM4] M. Goresky, R. Kottwitz, R. Macpherson, *Codimensions of root valuation strata*, Pure. Appl. Math. Q. 5 (2009), no. 4, 1253-1310.

[KL] D. Kazhdan, G. Lusztig, *Fixed point varieties on affine flag manifolds*, Israel. J. Math. **62**(1988), 129-168.

[L] G. Laumon, *Fibres de Springer et jacobiennes compactifiées*, Algebraic geometry and number theory, 515–563, Progr. Math., 253, Birkhäuser Boston, Boston, MA, 2006.

[Lu] V. Lucarelli, *Affine pavings for affine Springer fibers for split elements in* PGL_3, `http://arxiv.org/abs/math/0309132`

[Weil II] P. Deligne, *La conjecture de Weil, II*, Publ. Math. IHES, 52, 1980, 137-252.

EPFL SB MATHGEOM/GEOM, MA B1 447, STATION 8, CH-1015, LAUSANNE, SWITZERLAND
E-mail address: `zongbin.chen@gmail.com`